

IN THIS ISSUE:


1. Washoe County Quarterly Communicable Disease Statistics Q3 2018
2. Figure 1. Selected Notifiable Disease Report Trend.

**Reported Cases of Selected Communicable Disease
Washoe County, July – September 2018**

DISEASE	3 rd Quarter			Year to Date (Cumulative)		
	2016	2017	2018	2016	2017	2018
AIDS	3	6	4	9	10	10
Campylobacteriosis	15	8	15	35	29	32
Chlamydia	554	669	721	1693	1879	2031
Cryptosporidiosis	3	15	6	8	27	12
E. coli O157:H7, EHEC/STEC, HUS	0,2,0	2,4,2	1,3,1	1,2,0	2,6,2	2,9,1
Giardiasis	4	3	6	14	7	13
Gonorrhea	164	225	264	452	627	666
Haemophilus influenzae type b (Hib)	0	0	0	0	0	0
Hepatitis A (acute)	0	0	0	0	1	1
Hepatitis B (acute)	1	1	0	2	4	1
Hepatitis B (chronic)	21	14	8	48	48	44
Hepatitis C (acute)	3	3	0	4	12	3
Hepatitis C (chronic)	217	226	163	492	610	649
HIV	8	5	9	23	15	20
Influenza (Types A, B, & unknown)	13	39	12	1515**	1121	1570
Measles (confirmed)	0	0	0	0	0	1
Meningitis, Viral or Aseptic	4	28*	6	6	32*	17
Meningococcal Disease	0	0	0	0	0	0
Pertussis (confirmed & Probable)	0	2	3	0	7	9
Pneumococcal Disease, Invasive	10	10	8	45	41	48
Rabies (bat)	7	5	4	7	5	5
Rotavirus	7	0	2	12	10	9
RSV	2	3	4	367	579	390
Salmonellosis	11	12	15	23	23	32
Shigellosis	10*	1	3	12	2	5
Syphilis (Primary & Secondary)	8	19	33***	23	34	83***
Tuberculosis	1	1	3	3	11	7
West Nile Virus	1	23*	2	1	23*	2

* Outbreaks associated **During the several weeks in Q1, influenza like illness (ILI) activities were well beyond the region 9 baseline. ***Significantly increased morbidity.

FIGURE I. Selected notifiable disease reports, Washoe County, comparison of provisional 4-weeks totals , Week 36-39 (Ending on September29), 2018, with historical data



TO READ FIGURE 1

Figure 1 allows readers to compare reported morbidity for selected communicable diseases during the most current four weeks in comparison to the historical morbidities reported in 15 4-week periods in the past five years. For example, the total number of cases for P&S syphilis in the current 4 week period (week 36-39) was 13; however, the average of 4-week total for comparison periods during the past five years was only 3. The ratio of current 4-week total to mean of 15 4-week totals is 4.3, which means that the reported P&S syphilis is 4.3 times the average reported during the comparable time periods in the past five years.