

Community-wide Surveillance for Carbapenemase Producing Organisms (CPO) Statistical Report for 2024 Quarter 1

Division of Epidemiology & Public Health Preparedness (EPHP) 775-328-2447

Christabell Sotelo, MPH, Epidemiologist, Healthcare Associated Infection Coordinator,
csotelo@nnph.org

Cumulative Summary & Changes from Previous Quarter *

- CRO counts: 12
- CPO counts: 0
- CRO antibiotic resistance:
 - 3+ classes of antibiotics: 75.0%
 - 4+ classes of antibiotics: 58.3%
 - Pan resistance: 0

Please note caution should be taken when comparing 2023 and onward data to previous years as case definition changes have affected case counts.

*For definition and specifics on metrics summarized, please refer to corresponding sections and the surveillance definitions at the end.

CRO Overview

Table 1: CRO cases reported by quarter, Washoe County, 2024

CRO Type	Q1	Q2	Q3	Q4	Total
CRE	6	-	-	-	6
CRPA	6	-	-	-	6
CRA	0	-	-	-	0
Unknown	0	-	-	-	0
Other CROs	0	-	-	-	0
Total	12	-	-	-	12

For the current reporting quarter, 12 CROs were reported.

- 6 CREs, 6 CRPAs.

Table 1-1: Descriptive statistics for reported CRO cases, Washoe County, 2024


Characteristics		2024	
		No.	Percent (%)
Age	Median	77 years	NA
	Minimum	59 years	NA
	Maximum	91 years	NA
Gender	Male	5	41.67%
	Female	7	58.33%
Race/Ethnicity	White, non-Hispanic	12	100.00%
	White, Hispanic	0	0.00%
	Asian	0	0.00%
	Black	0	0.00%
	American Indian/Alaskan Native	0	0.00%
	Other	0	0.00%
	Unknown	0	0.00%
Washoe County Resident	Yes	11	91.67%
	No	1	8.33%
	Unknown	0	0.00%
Specimen Type	Urine	7	58.33%
	Respiratory	1	8.33%
	Wound	1	8.33%
	Rectal	0	0.00%
	Invasive (e.g., blood, cerebrospinal fluid)	0	0.00%
	Other	2	16.67%
	Surgical	1	8.33%
	Unknown	0	0.00%
Facility Type	Inpatient	5	41.67%
	Outpatient	3	25.00%
	Long Term Acute Care	0	0.00%
	Intensive Care Unit	3	25.00%
	Skilled Nursing Facility	1	8.33%
Total		12	100.00

In summary, 2024 CRO cases were:

- 77 years (median age).
- White, non-Hispanic (100%).
- Washoe County residents (91.67%).
- Detected from urine specimens (58.33%).
- Identified while in an inpatient facility (41.67%).

Data presented in this report is preliminary and may be updated in future reports as additional information is received throughout the year.

Figure 1: CRO cases reported by month, Washoe County, 2024



- The number of CRE cases were highest in January (4 reported) and lowest in February (1 reported) and March (1 reported).
- The number of CRPA were highest in January (3 reported) and lowest in March (1 reported).


Table 2: Proportion of CROs that were CREs, Washoe County, 2020-2024

Year	CRO Total	CRE Total	Proportion (%)
2020	90	48	53.33
2021	77	36	46.75
2022	145	62	42.76
2023	81	42	51.85
2024	12	6	50.00

- Of the 12 CRO’s reported, 51.85% (6/12) were CREs.

Data presented in this report is preliminary and may be updated in future reports as additional information is received throughout the year.

Figure 2: CRE cases reported by organism (N=6), Washoe County, 2024



EC-*Enterobacter cloacae*, ECo-*Escherichia coli*, SM-*Serratia marcescens*

Of the 6 CREs reported,

- *Enterobacter cloacae* was the most reported organism (67%).
- *Escherichia coli* was the least reported (16%).

Carbapenemase Producing Organisms (CPO)

Table 3: CPO cases reported, Washoe County, 2024

Month/ Year Reported	Resistance Mechanism	Organism	Clinical, Screening	Case notes
No CPOs reported in Q1.				

No CPOs were reported in quarter 1.

Tables 4 and 5 and Figures 3 and 4 present laboratory test data used to identify CPOs. The modified carbapenem inactivation method (mCIM) is a phenotypic (observable trait) test, while polymerase chain reaction (PCR) is a molecular test for carbapenemase genes. Please note the following when interpreting the data:

- Not all specimens are forwarded to the Nevada State Public Health Laboratory for testing.
- Some area hospitals perform PCR testing in-house.
- Though mCIM and PCR positive counts often match, in some instances, a specimen may only test positive for one of either test.

Data presented in this report is preliminary and may be updated in future reports as additional information is received throughout the year.


Table 4: Modified carbapenem inactivation method (mCIM) testing, Washoe County, 2020-2024

Year	N Tested	N Positive	Positivity (%)
2020	81	5	6.17
2021	71	5	7.04
2022*	109	3	2.75
2023	68	2	2.94
2024	9	0	0.00
Total	338	15	4.44

* One CPO is not included in Table 6 as they were identified by PCR testing and not mCIM.

- Out of the 9 specimens submitted for mCIM testing, none were positive (0/9).

Figure 3: Percent mCIM positive by organism, Washoe County, 2020-2024



KP-*Klebsiella pneumoniae*, PA-*Pseudomonas aeruginosa*, PP/PR-*Pseudomonas fluorescens/putida*, EC-*Enterobacter cloacae*, ECo- *Escherichia coli*

- In 2024, no organisms were mCIM positive (0%).
- From 2020-2023, the organisms that were mCIM positive varied.


Data presented in this report is preliminary and may be updated in future reports as additional information is received throughout the year.

Table 5: Polymerase chain reaction (PCR) testing, Washoe County, 2020-2024

Year	N Tested	N Positive	Positivity (%)
2020	7	5	71.4
2021	6	3	50.0
2022	6	4	66.7
2023	38	5	13.2
2024	5	0	0.0
Total	62	17	27.4

- Out of the 5 specimens submitted for PCR testing in 2024, none were positive (0/5).

Figure 4: Percent PCR positive by organism, Washoe County, 2020-2024



KP-*Klebsiella pneumoniae*, PA-*Pseudomonas aeruginosa*, PP/PR-*Pseudomonas fluorescens/putida*, EC-*Enterobacter cloacae*, KO-*Klebsiella oxytoca*


*1 screening specimen was PCR positive, but failed to culture an organism.

- In 2024, no organisms were PCR positive (0/5).
- From 2020-2023, the organisms that were PCR testing varied, however *Klebsiella pneumoniae* was the highest across all years combined.

Data presented in this report is preliminary and may be updated in future reports as additional information is received throughout the year.

Severity of Antibiotic Resistance

Figure 5: Percent of CRO cases resistant to classes of antibiotics, Washoe County, 2024



* Caution should be taken when comparing 2023 data to previous year as case definition change affected case counts.

In 2024, the proportion of reported CROs resistant to at least

- three or more classes of antibiotics was 75.0% (9/12).
- four or more classes of antibiotics was 58.3% (7/12).

Between 2020-2023, antibiotic resistance had a downward trend.

Table 6: Pan-resistance rate, Washoe County, 2020-2024

Year	Total N Cases	No. Pan-resistance	Proportion (%)	Organisms (No. pan-resistant)
2020	89	2	2.25	<i>Citrobacter sp.</i> (1), <i>K. pneumoniae</i> (1)
2021	76	0	0.00	-
2022	145	1	0.69	<i>Pseudomonas aeruginosa</i>
2023	81	1	1.23	<i>Acinetobacter baumannii</i>
2024	12	0	0.00	-

- Proportion pan-resistant*: 0% (0/12).

*Pan-resistance is defined as non-susceptible to all tested drugs at the clinical lab.

Data presented in this report is preliminary and may be updated in future reports as additional information is received throughout the year.

Table 7. Antibiotic Susceptibility for CRE, CRPA 2024

Antimicrobial Class or Subclass	CRE (n=6)			CRPA (n=6)		
	# Tested	# Susceptible	% Susceptible	# Tested	# Susceptible	% Susceptible
Penicillins						
Ampicillin	8	0	0.00	5	0	0.00
Piperacillin	-	-	-	1	0	0.00
Cephems						
Cefazolin	10	1	10.00	-	-	-
Cefepime	9	5	55.56	8	4	50.00
Ceftazidime	6	0	0.00	7	3	42.86
Ceftriaxone	10	1	10.00	-	-	-
Cefuroxime	4	1	25.00	-	-	-
β-Lactam/β-lactamase inhibitor combinations						
Amoxicillin-clavulanic acid	5	0	0.00	-	-	-
Ampicillin-sulbactam	8	0	0.00	5	0	0.00
Piperacillin-tazobactam	10	2	20.00	8	4	50.00
Ticarcillin-clavulanic acid	-	-	-	1	1	1.00
Fluoroquinolones						
Ciprofloxacin	10	8	80.00	10	4	40.00
Levofloxacin	10	8	80.00	6	2	33.33
Moxifloxacin	1	1	100.00	-	-	0.00
Aminoglycosides						
Amikacin	6	6	100.00	9	9	100.00
Gentamicin	10	10	100.00	9	7	77.78
Tobramycin	10	10	100.00	8	8	100.00
Sulfonamides						
Trimethoprim-sulfamethoxazole	10	9	90.00	-	-	-
Monobactams						
Aztreonam	4	0	0.00	6	3	50.00

Data presented in this report is preliminary and may be updated in future reports as additional information is received throughout the year.

Antimicrobial Class or Subclass	CRE (n=6)			CRPA (n=6)		
	# Tested	# Susceptible	% Susceptible	# Tested	# Susceptible	% Susceptible
Tetracyclines						
Tetracycline	6	4	66.67	-	-	-
Tigecycline	3	3	100.00	-	-	-
Nitrofurans						
Nitrofurantoin	4	3	75.00	-	-	-
Carbapenems						
Imipenem	1	1	0.00	5	0	0.00
Meropenem	7	6	85.71	10	5	50.00
Ertapenem	9	1	11.11			0.00

Surveillance Definitions (Years Updated)

Report Date (2024)

For this report, the date of specimen collection is used for case counts by months.

Carbapenemase-Producing Organisms (CPO) (2023)

Any specimen that meets confirmatory laboratory evidence:

- Positive phenotypic test for carbapenemase production **OR**
- Molecular test detecting a carbapenemase gene **OR**
- Next generation sequencing detecting a carbapenemase gene.

CPO cases will be classified as either clinical case (collected for diagnosing/treating disease), or as screening case (collected for detecting colonization), however since reason for collecting specimens is not reported, the specimen site denotes CPO case classification. Typically a CPO identified through a rectal, peri-rectal, axilla, groin, or stool specimen would be considered screening.

Duplicates (2023)

Duplicates are defined as the same organism/carbapenemase combination regardless of collection source and date. A screening case can be counted as a new clinical case if, for example, they developed a blood stream infection, found to be due to the same organism/carbapenemase combination, but a clinical case cannot be counted as a new screening case with same organism/carbapenemase combination.

Carbapenem Resistant *Enterobacteriaceae* (CRE) (2022)

Data presented in this report is preliminary and may be updated in future reports as additional information is received throughout the year.

Enterobacteriaceae that meets the following criteria:

- Resistant to ANY carbapenem antimicrobial (i.e., MIC of ≥ 4 mcg/ml for doripenem, meropenem, or imipenem OR ≥ 2 mcg/ml for ertapenem) **OR**
- Documented to produce carbapenemase

In addition:

- For bacteria that have intrinsic imipenem nonsusceptibility (i.e., *Morganella morganii*, *Proteus spp.*, *Providencia spp.*), resistant to carbapenems other than imipenem is required.

Carbapenem Resistant *Pseudomonas aeruginosa* (CRPA) (2022)

Pseudomonas aeruginosa isolated from any body site* that meets the following criteria:

- Resistant to imipenem, meropenem, or doripenem based on current Clinical and Laboratory Standards Institutes Standards (CLSI) M100 standards (≥ 8 mcg/mL); **AND/OR**
- Demonstrates production of a carbapenemase by a recognized method (e.g., CarbaNP or Polymerase chain reaction (PCR) or other methods).

*Excluding isolates from patients with cystic fibrosis (CF).

Carbapenem Resistant *Acinetobacter* (CRA) (2022)

Acinetobacter isolated from any body site that meets the following criteria:

- Resistant to imipenem, meropenem, or doripenem based on current Clinical and Laboratory Standards Institutes Standards (CLSI) M100 standards (≥ 8 mcg/mL); **AND/OR**
- Demonstrates production of a carbapenemase by a recognized method (e.g., CarbaNP or PCR or other methods).

Carbapenem Resistant Organisms (CRO) (2017)

Any organisms meeting the above definitions for CRE, CRPA, and CRA are considered CRO.

Carbapenemase Producing Organisms (CPO) (2017)

Any organisms producing carbapenemase which is laboratory-confirmed are defined as CPO.

Multi-Drug Resistant Bacilli – Carbapenem Resistant (MDRB-CR) (2010-2016)

A case is defined as an infection with an MDRB-CR organism of one patient per hospitalization per year regardless of resident status. Infection with a second species of MDRB-CR organism in the same patient is counted as a separate case. Infections with those Gram-negative bacilli that are constitutively resistant to carbapenems, specifically *Stenotrophomonas*, *Aeromonas* & *Chryseobacterium*, are not counted as cases.

Data presented in this report is preliminary and may be updated in future reports as additional information is received throughout the year.

MDRB-CR organisms refer to Gram negative bacilli that are resistant to three or more classes of antibiotics, one of which must be Carbapenem.

Patient's Residency (SINCE 2010)

Patients from out of jurisdiction (OOJ) are included in the surveillance report as long as isolates meet the above surveillance definitions.